SUBJECT : CHEMISTRY	DAY-2		
SESSION : AFTERNOON	TIME: 02.30 P.M. TO 03.50 P.M.		

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
60	80 MINUTES	70 MINUTES

QUESTION BOOKLET DETAILS			
VERSION CODE	SERIAL NUMBER		
A - 1	807425		
	VERSION CODE		

## DOs:

- 1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
- This Question Booklet is issued to you by the invigilator after the 2<sup>nd</sup> Bell i.e., after 02.30 p.m.
- The Serial Number of this question booklet should be entered on the OMR answer sheet.
- The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
- Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

## DON'TS:

- THE TIMING AND MARKS PRINTED ON THE OMR ANSWER SHEET SHOULD NOT BE DAMAGED/MUTILATED/SPOILED.
- The 3<sup>rd</sup> Bell rings at 02.40 p.m., till then;
  - Do not remove the paper seals present on all the 3 sides of this question booklet.
  - Do not look inside this question booklet.
  - Do not start answering on the OMR answer sheet.

## IMPORTANT INSTRUCTIONS TO CANDIDATES

- This question booklet contains 60 questions and each question will have one statement and four distracters. (Four different options / choices.)
- After the 3<sup>rd</sup> Bell is rung at 02.40 p.m., remove the paper seals of this question booklet and check that this
  booklet does not have any unprinted or torn or missing pages or items etc., if so, get it replaced by a complete test
  booklet. Read each item and start answering on the OMR answer sheet.
- During the subsequent 70 minutes:
  - Read each question carefully.
  - Choose the correct answer from out of the four available distracters (options / choices) given under each question / statement.
  - Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALL POINT PEN
    against the question number on the OMR answer sheet.

## CORRECT METHOD OF SHADING THE CIRCLE ON THE OMR SHEET IS AS SHOWN BELOW:

- Please note that even a minute unintended ink dot on the OMR answer sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
- After the last bell is rung at 03.50 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 7. Hand over the OMR ANSWER SHEET to the room invigilator as it is.
- After separating the top sheet (Our Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- Preserve the replica of the OMR answer sheet for a minimum period of ONE year.

C

Turn Over

- The mass of a non-volatile solute of molar mass 40 g mol<sup>-1</sup> that should be dissolved in 114 g of octane to lower its vapour pressure by 20% is
  - (1) 10 g

(2) 11.4 g

(3) 9.8 g

- (4) 12.8 g
- 2. During the adsorption of a gas on the surface of a solid, which of the following is TRUE?
  - (1)  $\Delta G < O, \Delta H > O, \Delta S < O$
  - (2) ΔG > O, ΔH < O, ΔS < O</p>
  - (3) ΔG < O, ΔH < O, ΔS < O</p>
  - (4)  $\Delta G < O$ ,  $\Delta H < O$ ,  $\Delta S > O$
- The approximate time duration in hours to electroplate 30g of calcium from molten calcium chloride using a current of 5 amp is

[At. mass of Ca = 40]

(1) 8

(2) 80

(3) 10

- (4) 16
- 4. The pH of the solution obtained by mixing 100 ml of a solution of pH = 3 with 400 ml of a solution of pH = 4 is
  - (1) 3 log 2.8

(2)  $7 - \log 2.8$ 

(3) 4 - log 2.8

(4) 5 - log 2.8

5. The equilibrium constant of the reaction :

$$A_{(s)} + 2B_{(aq)}^+ \Longrightarrow A_{(aq)}^{2+} + 2B_{(s)}; E_{cell}^o = 0.0295 \text{ V} \text{ is}$$

$$\left[\frac{2.303 \text{ RT}}{\text{F}} = 0.059\right]$$

(1) 10

(2)  $2 \times 10^2$ 

(3) 3 × 10<sup>2</sup>

(4) 2 × 10<sup>5</sup>

6. An oxygen containing organic compound was found to contain 52% carbon and 13% of hydrogen. Its vapour density is 23. The compound reacts with sodium metal to liberate hydrogen. A functional isomer of this compound is

(1) Ethanol

(2) Ethanal

(3) Methoxy Methane

(4) Methoxy Ethane

7. Which one of the following is NOT true regarding electromeric effect?

(1) It results in the appearance of partial charges on the carbon atoms.

(2) It is a temporary effect.

(3) It operates on multiple bonds.

(4) It requires an attacking reagent.

8. Which one of the following is NOT formed when a mixture of methyl bromide and bromobenzene is heated with sodium metal in the presence of dry Ether?

(1) Ethane

(2) Diphenyl

(3) Propane

(4) Toluene

(3) 80% Ethanol + 20% Benzene + Small quantity of Petrol

(4) 50% Petrol + 50% Ethanol + Small quantity of Benzene

10. Identify 'C' in the following:

$$\bigcirc + CH_3 - CH - CH_3 \xrightarrow{Anhyd \ A/Cl_3} A \xrightarrow{O_2} B \xrightarrow{dil \ H_2SO_4} Phenol + C$$

(1) Water

(2) Ethanol

(3) Propanone

(4) Cumene hydroperoxide

 20 ml of methane is completely burnt using 50 ml of oxygen. The volume of the gas left after cooling to room temperature is

(1) 80 ml

(2) 40 ml

(3) 60 ml

(4) 30 ml

12. 100 ml of 0.1 M acetic acid is completely neutralized using a standard solution of NaOH. The volume of Ethane obtained at STP after the complete electrolysis of the resulting solution is

(1) 112 ml

(2) 56 ml

(3) 224 ml

(4) 560 ml

	(1)	Cellulose	(2)	Toluene		
	(3)	Cyclohexane	(4)	Starch		
14.	Which of	the following is NO	T TRUE for S <sub>N</sub> 1 react	ion?		
	(1)	Favoured by polar	solvents.			
	(2)	3° – alkyl halides §	generally react through	S <sub>N</sub> 1 reaction.		
	(3)	The rate of the re nucleophile.	action does not deper	nd upon the mola	ar concent	ration of the
	(4)	1° – alkyl halides g	generally react through	S <sub>N</sub> 1 reaction.		
15.	Oil of wi	nter green is				
	(1)	an ester	(2)	a carboxylic aci	d	
	(3)	an alcohol	(4)	a ketone		
16.			rns with a sooty flame		wards Tol	len's reagent
	test and p	ositive for Borsche's	s reagent test. The con	ipound 'A' is		
	(1)	Benzaldehyde	(2)	Acetophenone		
	(3)	Acetone	(4)	Salicylic acid		
			Space For Rough We	ork		

13. Saccharin, an artificial sweetner, is manufactured from

17. For a reaction: A + B → Products, the rate of the reaction at various concentrations are given below:

Expt No	[A]	[B]	rate (mol dm <sup>-3</sup> s <sup>-1</sup> )
1	0.2	0.2	2
2	0.2	0.4	4
3	0.6	0.4	36

The rate law for the above reaction is

(1)  $r = K[A]^2[B]$ 

(2) r = K[A][B]<sup>2</sup>

(3)  $r = K[A]^3[B]$ 

(4)  $r = K[A]^2[B]^2$ 

18. Which one of the following has NO unpaired electrons?

(1) O<sub>2</sub>

(2) 0

(3) O<sub>2</sub><sup>+</sup>

(4) O<sub>2</sub>

19. The atomic number of cobalt is 27. The EAN of cobalt in Na<sub>3</sub>[Co(NO<sub>2</sub>)<sub>4</sub>Cl<sub>2</sub>] is

(1) 35

(2) 24

(3) 36

(4) 34

20. The "spin only" magnetic moment of Ni2+ in aqueous solution would be

[At No. of Ni = 28]

(1) √6 BM

(2) √15 BM

(3)  $\sqrt{2}$  BM

(4) √8 BM

21.	Impossib	le orbital among the following is		
	(1)	2s	(2)	3f
	(3)	2p	(4)	4d
22.	The total	number of electrons in 18 ml of v	water (der	nsity = $1 \text{ g ml}^{-1}$ ) is
	(1)	$6.02 \times 10^{23}$	(2)	$6.02 \times 10^{25}$
	(3)	$6.02 \times 10^{24}$	(4)	$6.02 \times 18 \times 10^{23}$
23.	The num	ber of moles of hydrogen that can	be added	d to 1 mole of an oil is the highest in
	(1)	Linseed oil	(2)	Groundnut oil
	(3)	Sunflower seed oil	(4)	Mustard oil
24.	The react	ion between sodium and water ca	in be mad	le less vigorous by
	(1)	lowering the temperature	(2)	adding a little alcohol
	(3)	amalgamating sodium	(4)	adding a little acetic acid
25.	All collo	idal dispersions have		
	(1)	very high osmotic pressure	(2)	low osmotic pressure
	(3)	no osmotic pressure	(4)	high osmotic pressure
26.	Silver ioc	lide is used for producing artificia	ıl rain bec	cause AgI
	(1)	is easy to spray at high altitude		
	(2)	is easy to synthesize		
	(3)	has crystal structure similar to it	ce	
	(4)	is insoluble in water		

27. The equilibrium constant of a reaction is 0.008 at 298 K. The standard free energy change of the reaction at the same temperature is

(1) +11.96 kJ

(2) -11.96 kJ

(3) -5.43 kJ

(4) -8.46 kJ

28. The function of potassium ethyl xanthate in froth floatation process is to make the ore

- (1) attracted towards water
- (2) water repellant

(3) lighter

(4) heavier

29. The correct order of electronegativities of N, O, F & P is

(1) F>N>P>O

(2) F > O > P > N

(3) F>O>N>P

(4) N>O>F>P

30. The s-block element used as a catalyst in the manufacture of Buna-S rubber is

(1) Mg

(2) C:

(3) Ba

(4) Na

31. Which of the following is NOT a characteristic of a covalent compound?

- (1) Low melting point
- (2) No definite geometry
- (3) Insoluble in polar solvent
- (4) Small difference in electronegativity between the combining atoms.

	(1)	125 ml		(2)	25 ml	
	(3)	12.5 ml		(4)	37.5 ml	
		saleden von				
33.	A ligand					
	(1)	Lewis acid				
	(2)	Bronsted acid				
	(3)	either a Lewis acid	d or a Lewis bas	se		
	(4)	Lewis base				
34.					neir pure states are in the ratio	
34.	A binary		B contains A an	nd B in	neir pure states are in the ratio the mole proportion of 1:2.7	
34.	A binary	solution of A and	B contains A an	nd B in	neir pure states are in the ratio the mole proportion of 1:2.7	
34.	A binary fraction o	solution of A and of A in the vapour p	B contains A an	nd B in tion wil (2)	neir pure states are in the ratio the mole proportion of 1:2.7	
	A binary fraction of (1) (3)	solution of A and of A in the vapour p 0.33	B contains A ar hase of the solut	nd B in tion wil (2) (4)	neir pure states are in the ratio the mole proportion of 1 : 2. I be	
	A binary fraction of (1) (3)	solution of A and of A in the vapour p 0.33 0.25	B contains A an hase of the solut ments is TRUE	nd B in tion wil (2) (4)	neir pure states are in the ratio the mole proportion of 1 : 2. 7 lbe  0.2  0.52	
	A binary fraction of (1) (3) Which of	solution of A and of A in the vapour plus 0.33 0.25 the following state.  The total entropy	B contains A ar hase of the solut ments is TRUE	(2) (4) ?	neir pure states are in the ratio the mole proportion of 1 : 2. 7 lbe  0.2  0.52	
	A binary fraction of (1) (3) Which of (1)	solution of A and of A in the vapour plus 0.33 0.25  The following states The total entropy The total entropy	B contains A ar hase of the solut ments is TRUE of the universe i	nd B in tion wil (2) (4) ? remains is contin	neir pure states are in the ratio the mole proportion of 1 : 2. 7 lbe  0.2  0.52  constant.	

- 5 ml of 0.4 N NaOH is mixed with 20 ml of 0.1 N HCl. The pH of the resulting solution will be
  - (1) 6

(2) 7

(3) 8

- (4) 5
- 37. On adding which of the following, the pH of 20 ml of 0.1 N HCl will not alter?
  - (1) 1 ml of 1 N HCI

- (2) 20 ml of distilled water
- (3) 1 ml of 0.1 N NaOH
- (4) 500 ml of HC l of pH = 1
- 38. Which one of the following has a potential more than zero?
  - (1) Pt,  $\frac{1}{2}$  H<sub>2</sub> (1 atm) | HCl (1 M)
  - (2) Pt,  $\frac{1}{2}$  H<sub>2</sub> (1 atm) | HCl (2 M)
  - (3) Pt,  $\frac{1}{2}$  H<sub>2</sub> (1 atm) | HCI (0.1 M)
  - (4) Pt,  $\frac{1}{2}$  H<sub>2</sub> (1 atm) | HCI (0.5 M)
- HCHO was treated with a reagent X. The product formed upon hydrolysis in the presence of an acid gave C<sub>2</sub>H<sub>5</sub>OH. The reagent X is
  - (1) aqueous KOH

(2) alcoholic KOH

(3) alcoholic KCN

(4) CH<sub>3</sub> MgI

40. Benzylamine is a stronger base than aniline because

The lone pair of electrons on the nitrogen atom in benzylamine is delocalised.

The lone pair of electrons on the nitrogen atom in aniline is delocalised. (2)

The lone pair of electrons on the nitrogen atom in aniline is not involved in (3) resonance.

Benzylamine has a higher molecular mass than aniline.

41. The relative acidic strengths of benzoic acid, o-toluic acid and p-toluic acid is of the decreasing order:

> p-toluic acid > o-toluic acid > benzoic acid (1)

> (2) o-toluic acid > p-toluic acid > benzoic acid

> p-toluic acid > benzoic acid > o-toluic acid (3)

> o-toluic acid > benzoic acid > p-toluic acid


42. The C-H bond and C-C bond in ethane are formed by which of the following types of overlap?

(1) 
$$sp^3 - s$$
 and  $sp^3 - sp^3$ 

(2) 
$$sp^2 - s$$
 and  $sp^2 - sp^2$ 

(3) 
$$sp - s$$
 and  $sp - sp$ 

43. The IUPAC name of



4-Hydroxy-2-pentanone (2) 2-Hydroxy-4-pentanone

(3) 2-Oxo-4-pentanol

(4) 4-Keto-2-pentanol

44. A first order reaction is 60% complete in 20 minutes. How long will the reaction take to be 84% complete?

(1) 54 mins

(2) 68 mins

(3) 40 mins

(4) 76 mins

45. A given sample of milk turns sour at room temperature (27 °C) in 5 hours. In a refrigerator at -3 °C, it can be stored 10 times longer. The energy of activation for the souring of milk is

(2) 2.303 × 5 R kJ · mol<sup>-1</sup>

(4) 2.303 × 2.7 R kJ - mol<sup>-1</sup>

46. At 300 K, a gaseous reaction:

$$A \rightarrow B + C$$

was found to follow first order kinetics. Starting with pure A, the total pressure at the end of 20 minutes was 100 mm of Hg. The total pressure after the completion of the reaction is 180 mm of Hg. The partial pressure of A (in mm of Hg) is

(1) 100

(2) 90

(3) 180

(4) 80

47. From the Ellingham graphs on carbon, which of the following statements is FALSE?

- CO<sub>2</sub> is more stable than CO at less than 983 K
- (2) CO reduces Fe<sub>2</sub>O<sub>3</sub> to Fe at less than 983 K
- (3) CO is less stable than CO<sub>2</sub> at more than 983 K
- (4) CO reduces Fe2O3 to Fe in the reduction zone of Blast furnace

- 48. Which of the following is a negatively charged bidentate ligand?
  - Dimethyl glyoximato
- Cyano

Ethylene diamine

- Acetato
- 49. The secondary valency of platinum in tetra ammine dichloroplatinum (IV) chloride is
  - (1) +4

(2) +2

(3)

- (4) 6
- 50. Which one of the following has a magnetic moment of 1.75 BM?
- (1) Ti<sup>3+</sup> (2) V<sup>3+</sup>

- (3) Cr<sup>3+</sup> (4) Fe<sup>3+</sup>
- 51. The correct order of ionisation energy of C, N, O & F is
  - (1) F < N < C < O

(2) C < N < O < F

(3) C < O < N < F

- (4) F < O < N < C
- 52. The correct set of four quantum numbers for the outermost electron of sodium (Z = 11) is
  - (1) 3, 1, 0,  $\frac{1}{2}$

(2)  $3, 1, 1, \frac{1}{2}$ 

(3) 3, 2, 1,  $\frac{1}{2}$ 

(4)  $3, 0, 0, \frac{1}{2}$ 

54.	The equi chloride		certain bivalent	metal is 2	20. The molecular mass of its anhydrous
	(1)	91		(2)	111
	(3)	55.5		(4)	75.5
55.					t 298 K and under 1 atm pressure. It is omposes to NO <sub>2</sub> . The resulting pressure
	(1)	2.4 atm		(2)	1.2 atm
	(3)	4.8 atm		(4)	2.8 atm
56.	Sucrose i	s NOT a reducing	sugar since		
	(1)	it is chemically	stable		
	(2)	it contains no fre	ce aldehyde or k	ceto group	adjacent to a CHOH group
	(3)	it is built up of a	fructose unit		
	(4)	it is optically act	tive		
-			Space For I	Rongh Wo	ork

Cryolite

Calamine

(2)

(4)

53. The ore that is concentrated by the Froth Floatation process is

(1)

(3)

Chalcopyrites

Cuprite

	(3)	IVACIV		(4)	Naire		
58.	Dialysis	can be used to s	eparate				
	(1)	glucose & fru	ctose	(2)	protein & starch		
	(3)	glucose & pro	tein	(4)	glucose & NaCl		
59.	The percentage respective		haracter of the	hybrid o	rbitals in graphit	e and d	iamond ar
	(1)	33 and 25		(2)	50 and 75		
	(3)	67 and 75		(4)	33 and 75		
		1.1					
60.	A gas ex	pands from a vo	olume of 1 m <sup>3</sup> to	a volume	of 2 m3 against ar	n external	pressure o
	10 <sup>5</sup> Nm <sup>-2</sup>	2. The work don	e by the gas will	be			
	(1)	$10^5  \mathrm{kJ}$		(2)	$10^2  \mathrm{kJ}$		
	(3)	$10^2\mathrm{J}$		(4)	$10^3  \mathrm{J}$		
-			Space For	Rough Wo	ork		

57. Which one of the following contains ionic, covalent and co-ordinate bonds?

(1) NaOH

NaCl

(2)



A-1

16

C

SEAL