SUBJECT : CHEMISTRY DAY-2

SESSION : AFTERNOON TIME : 02.30 P.M. TO 03.50 P.M.

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
60	80 MINUTES	70 MINUTES

MENTION YOUR	QUESTION BOOKLET DETAILS		
CET NUMBER	VERSION CODE	SERIAL NUMBER	
5- 8	A-1	729873	

DOs:

- 1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
- This Question Booklet is issued to you by the invigilator after the 2nd Bell i.e., after 2.30 p.m.
- 3. The Serial Number of this question booklet should be entered on the OMR answer sheet.
- The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
- Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

DON'TS:

- THE TIMING AND MARKS PRINTED ON THE OMR ANSWER SHEET SHOULD NOT BE DAMAGED / MUTILATED / SPOILED.
- The 3rd Bell rings at 2.40 p.m., till then;
 - Do not remove the paper seal present on the right hand side of this question booklet.
 - · Do not look inside this question booklet.
 - Do not start answering on the OMR answer sheet.

IMPORTANT INSTRUCTIONS TO CANDIDATES

- This question booklet contains 60 questions and each question will have one statement and four distracters. (Four different options / choices.)
- After the 3rd Bell is rung at 2.40 p.m., remove the paper seal on the right hand side of this question booklet and
 check that this booklet does not have any unprinted or torn or missing pages or items etc., if so, get it replaced by
 a complete test booklet. Read each item and start answering on the OMR answer sheet.
- 3. During the subsequent 70 minutes:
 - · Read each question carefully.
 - Choose the correct answer from out of the four available distracters (options / choices) given under each question / statement.
 - Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALL POINT PEN
 against the question number on the OMR answer sheet.

Correct Method of shading the circle on the OMR answer sheet is as shown below:



- Please note that even a minute unintended ink dot on the OMR answer sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
- Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
- After the last bell is rung at 3.50 p.m., stop writing on the OMR answer sheet and affix your LEFT HAND THUMB IMPRESSION on the OMR answer sheet as per the instructions.
- 7. Hand over the OMR ANSWER SHEET to the room invigilator as it is.
- After separating the top sheet (Our Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
- 9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.

C

Turn Over

- 1. The unit cell with crystallographic dimensions, $a \neq b \neq c$, $\alpha = \gamma = 90$ and $\beta \neq 90$ is
 - (1) Triclinic

(2) Monoclinic

(3) Orthorhombic

- (4) Tetragonal
- 2. While charging the lead storage battery.
 - (1) PbSO₄ on anode is reduced to Pb
 - (2) PbSO₄ on cathode is reduced to Pb
 - (3) PbSO4 on cathode is oxidized to Pb
 - (4) PbSO₄ on anode is oxidized to PbO₅
- 3. Adenosine is an example of
 - (1) Nucleotide

(2) Purine base

(3) Pyrimidine base

(4) Nucleoside

- 4. Orlon has monomeric unit
 - (1) Acrolein

(2) Glycol

(3) Vinyl cyanide

- (4) Isoprene
- 5. The two electrons have the following set of quantum numbers :

$$P = 3, 2, -2, +\frac{1}{2}$$

$$Q = 3, 0, 0, +\frac{1}{2}$$

Which of the following statement is true?

- (1) P and Q have same energy
- (2) P has greater energy than Q
- (3) P has lesser energy than Q
- (4) P and Q represent same electron

6.	LE (5	ienimi	tor	covi	dise
O.	II of	37	Сшп	IOL	OAI	UISC

(1) PbS

(2) Na₂SO₃

(3) O₃

(4) KI

7. In the given set of reactions,

2-Bromopropane
$$\xrightarrow{\text{AgCN}}$$
 X $\xrightarrow{\text{LiA/H}_4}$ Y

the IUPAC name of product 'Y' is

- (1) N-Methylpropanamine
- (2) N-Isopropylmethanamine

(3) Butan-2-amine

- (4) N-Methylpropan-2-amine
- 8. On heating with concentrated NaOH solution in an inert atmosphere of CO₂, white phosphorous gives a gas. Which of the following statement is <u>incorrect</u> about the gas?
 - It is less basic than NH₃.
 - (2) It is more basic than NH₃.
 - (3) It is highly poisonous and has smell like rotten fish.
 - (4) It's solution in water decomposes in the presence of light.
- Sodium metal crystallizes in B.C.C. lattice with edge length of 4.29 Å. The radius of sodium atom is
 - (1) 2.857 Å

(2) 1.601 Å

(3) 2.145 Å

C

(4) 1.857 Å

10.	0.06% (w	//v) aqueous solution of urea	is isotonic wit	th
	(1)	0.06% glucose solution	(2)	0.6% glucose solution
	(3)	0.01 M glucose solution	(4)	0.1 M glucose solution
11.		order reaction, the concentral is it half completed?	tion of the rea	actant is reduced to 12.5% in one hour.
	(1)	3 hr	(2)	20 min
	(3)	30 min	(4)	15 min
12.	The elect	rolyte having maximum floco	culation value	for Agl/Ag* sol. is
	(1)	NaC/	(2)	Na ₂ S
	(3)	Na ₂ SO ₄	(4)	Na ₃ PO ₄
13.		s extracted from Copper pyrion the principle that	tes by heating	g in a Bessemer converter. The method
	(1)	Copper has more affinity fo	r oxygen than	Sulphur at high temperature.
	(2)	Iron has less affinity for oxy	ygen than Sul	phur at high temperature.
	(3)	Copper has less affinity for	oxygen than	Sulphur at high temperature.
	(4)	Sulphur has less affinity for	oxygen at hi	gh temperature.
14.	Which of	f the following will be able to	show geome	trical isomerism?
	(1)	MA ₃ B – Square planar	(2)	MA ₂ B ₂ - Tetrahedral

Space For Rough Work

(3) MABCD - Square planar (4) MABCD - Tetrahedral

- 15. The electronic configuration of Gd2+ is (at. no. of Gd is 64)
 - (1) [Xe] 4f8

(2) [Xe] 4f

(3) [Xe] 4f7 5d1 6s2

- (4) [Xe] 4f7 5d1
- 16. $MSO_4 \xrightarrow{NH_4OH} \downarrow X \xrightarrow{NH_4OH} Y \xrightarrow{H_2S} \downarrow Z$

Here M and Z are

(1) Cu, ZnS

(2) Zn, ZnS

(3) Fe, FeS

- (4) Al, Al, S,
- 17. The hydrolysis of optically active 2-bromobutane with aqueous NaOH result in the formation of
 - (1) (+) butan-2-ol

(-) butan-2-ol (2)

(3) (±) butan-1-ol

- (4) (±) butan-2-ol
- 18. The distinguishing test between methanoic acid and ethanoic acid is
 - (1) Litmus test

Tollen's test


Esterification test (3)

- (4) Sodium bicarbonate test
- 19. In H2 O2 fuel cell the reaction occurring at cathode is


 - $(1) \quad 2H_{2(g)} + O_{2(g)} \longrightarrow 2H_2O_{(l)} \qquad (2) \quad O_{2(g)} + 2H_2O_{(l)} + 4e^- \longrightarrow 4\overline{O}H_{(aq)}$


 - (3) $H^+ + e^- \longrightarrow \frac{1}{2} H_2$ (4) $H^+_{(aq)} + \overline{O}H_{(aq)} \longrightarrow H_2O_{(l)}$

20. Which of the following curve is in accordance with Freundlich adsorption isotherm?









21. How many ions per molecule are produced in the solution when Mohr salt is dissolved in excess of water?

(1) 4

(2)

(3) 6

(4) 10

22. Glycogen is

- a polymer of β-D-glucose units
- (2) a structural polysaccharide
- (3) structurally very much similar to amylopectin
- (4) structurally similar to amylopectin but extensively branched

23. Number of possible alkynes with formula C5H8 is

(1) 2

(2) 3

(3) 4

(4) 5

24.	Which of	the following	aqueous solution	has the highes	t freezing point?
-----	----------	---------------	------------------	----------------	-------------------

(1) 0.1 M Sucrose

(2) 0.01 M NaC/

(3) 0.1 M NaC/

(4) 0.01 M Na₂SO₄

 Half life period of a first order reaction is 10 min. Starting with initial concentration 12 M, the rate after 20 min is

(1) 0.0693 M min-1

(2) 0.693 × 3 M min-1

(3) 0.0693 × 3 M min-1

(4) 0.0693 × 4 M min-1

26. The salt which responds to dilute and concentrated H2SO4 is

(1) CaF₂

(2) Ba(NO₃)₂

(3) Na₂SO₄

(4) Na₃PO₄

27. On heating potassium permanganate, one of the following compound is not obtained:

(1) O₂

(2) MnO

(3) MnO₂

(4) K2MnO4

28. $Br + Mg \xrightarrow{dry \text{ ether}} A \xrightarrow{H_2O} B$.

The product 'B' is

(1) OH

(2) — MgBr

(3)

(4) O OH

(1)	Nucl	eophilic	substitut	tion	(2)	Nucleophilic addition
(3)		and the later	addition		(4)	Electrophilic substitution
One of	the follo	wing is	an essent	ial amino	acid.	
(1)					(2)	Cysteine
(3)	Isole	ucine			(4)	Serine
The agu	eous so	ution of	followin	n salt will	have the	lowest pH:
(1)			101101111	ig sait will	(2)	NaC/O
(3)	NaC/	02			(4)	
For one below:						ion enthalpies (in kJ mol-1) are gi
For one below :	1 st	2 nd	3 rd	4 th	5 th	
I.E.	1 st 577.5					
I.E.	1 st 577.5 ment is	2 nd	3 rd	4 th	5 th 14,820	
I.E. The eler	1 st 577.5 ment is	2 nd	3 rd	4 th	5 th	P
I.E. The eler	1 st 577.5 nent is	2 nd	3 rd	4 th	5 th 14,820	P
1.E The eler (1) (3)	1 st 577.5 nent is Si Al f an org	2 nd 1810	3 rd 2750	4 th 11,580	5 th 14,820 (2) (4) C, H and	P
1.E The eler (1) (3) 0.30 g o CO ₂ and	1 st 577.5 ment is Si Al f an org 1 0.18 g nd is	2 nd 1810 anic con H ₂ O. If	3 rd 2750	4 th 11,580	5 th 14,820 (2) (4) C, H and	P Mg
The eler (1) (3) 0.30 g o CO ₂ and compou	1 st 577.5 ment is Si Al of an org	2 nd 1810 anic con H ₂ O. If	3 rd 2750	4 th 11,580	5 th 14,820 (2) (4) C, H and ound wei	P Mg d Oxygen on combustion yields 0.4 ghs 60, then molecular formula of

. Ch	eilosis	and digestive disor	rders are due to t	he defic	ciency of	
	(1)	Thiamine		(2)	Ascorbic acid	
	(3)	Riboflavin		(4)	Pyridoxine	
	w mar		ectricity are requ	iired fo	r the oxidation of one i	mol of water to
	(1)	$9.65 \times 10^{4} \text{ C}$		(2)	$1.93\times10^4\mathrm{C}$	
	(3)	$1.93 \times 10^5 \text{ C}$		(4)	19.3 × 10 ⁵ C	
7. 100) cm ³	The state of the s			of 2 M CH ₃ OH to for	
	inge in	the initial rate if ea	ach solution is di	iluted w	itin equal volume of wa	ter would be
	inge in	the initial rate if ea 2 times	ach solution is di	(2)	4 times	ter would be
			ach solution is di			ter would be
	(1)	2 times	Space For Re	(2) (4)	4 times 0.25 times	ter would be
	(1)	2 times		(2) (4)	4 times 0.25 times	ter would be
	(1)	2 times		(2) (4)	4 times 0.25 times	ter would be

A-1

34. One of the following amide will not undergo Hoffmann bromamide reaction:

(1)

(2)

(3)

C

CH3CONH2

CH3CONHCH3

C6H5CONH2

(4) CH3CH2CONH2

8.	Which of	f the following colloids cannot be	easily coa	igulated?
	(1)	Lyophobic colloids		
	(2)	Multimolecular colloids		
	(3)	Macromolecular colloids		
	(4)	Irreversible colloids		
39.	The com	plex ion having minimum magnit	tude of Δ_0	(CFSE) is
	(1)	[Cr(CN) ₆] ³	(2)	[Co(NH ₃) ₆] ³⁺
	(3)	$[Co(Cl)_6]^{3-}$	(4)	[Cr(H ₂ O) ₆] ³⁺
40.	The arra	ngement of following compounds		
	i. bro	momethane		
	ii. bro	moform		
	iii. chle	oromethane		
	iv. dib	romomethane		
	In the in	creasing order of their boiling po	int is	
	(1)	iii < i < iv < ii	(2)	iv < iii < i < ii
	(3)	ii < iii < i < iv	(4)	i < ii < iii < iv

41. Iodoform can be prepared from all, except


(1) propan-2-ol

(2) butan-2-one

(3) propan-1-ol

(4) acetophenone

42. Identify 'Q' in the following sequence of reactions:



- 43. Cryolite is
 - Na₃A/F₆ and is used in the electrolysis of alumina for decreasing electrical conductivity.
 - Na₃A/F₆ and is used in the electrolysis of alumina for lowering the melting point of alumina only.
 - (3) Na₃A/F₆ and is used in the electrolysis of alumina for lowering the melting point and increasing the conductivity of alumina.
 - (4) Na₃A/F₆ and is used in the electrolytic refining of alumina.

44. Which of the following compound of Xenon has pyramidal geometry?

(1) XeOF₄

(2) XeF

(3) XeO₃

(4) XeF₄

45. After adding non-volatile solute freezing point of water decreases to –0.186 °C. Calculate ΔT_b if K_f = 1.86 K kg mol $^{-1}$ and K_b = 0.521 K kg mol $^{-1}$


(1) 0.521

(2) 0.0521

(3) 1.86

(4) 0.0186

46. Plot of Maxwell's distribution of velocities is given below:



Which of the following is correct about this plot?

(1) $T_1 \le T_2$

(2) f₁ > f₂

(3) $T_1 > T_2$

(4) V₁ < V

47. The pair of compound which cannot exist together in solution is

- (1) NaHCO3 and NaOH
- (2) NaHCO3 and H2O
- (3) NaHCO3 and Na2CO3
- (4) Na2CO3 and NaOH

48. What amount of dioxygen (in gram) contains 1.8×10^{22} molecules?

(1) 0.0960

(2) 0.960

(3) 9.60

(4) 96.0

- Using MOT, compare O₂⁺ and O₂ species and choose the incorrect option.
 - (1) O2 have higher bond order than O2.
 - (2) O₂ is less stable.
 - (3) O₂⁺ is diamagnetic while O₂⁻ is paramagnetic.
 - (4) Both O₂ and O₂ are paramagnetic.
- 50. Which of the following is not true?
 - (1) Erythromycin is a bacteriostatic antibiotic.
 - (2) Ampicillin is not a natural antibiotic.
 - (3) Prontosil is not converted into sulphanilamide in the body.
 - (4) Vancomycin is a broad spectrum antibiotic.
- 51. In the reaction

$$S + \frac{3}{2}O_2 \longrightarrow SO_3 + 2v \text{ kJ} \text{ and } SO_2 + \frac{1}{2}O_2 \longrightarrow SO_3 + y \text{ kJ}$$

heat of formation of SO2 is

(1) x + y

(2) x-y

(3) 2x - y

- (4) 2x + y
- 52. Arrange the following compounds in the increasing order of their acidic strength:
 - i. m-nitrophenol
- ii. m-cresol

iii. phenol

- iv. m-chlorophenol
- (1) iii < ii < i < iv

(2) ii < iv < iii < i

(3) ii < iii < iv < i

(4) ii < iii < i < iv

53. In the sequence of following reactions:

$$P \xrightarrow{\text{(1) Br}_2} Q \xrightarrow{\text{(2) NaNO}_2/HC/} R \xrightarrow{\text{EMnO}_4} R$$

$$Q \xrightarrow{\text{(2) NaNO}_2/HC/} R \xrightarrow{\text{OH}} R$$

the starting compound 'P' is

(1) o-nitro toluene

(2) m-nitro toluene

(3) o-bromo toluene

(4) p-nitro toluene

 Acetic acid is treated with Ca(OH)₂ and the product so obtained is subjected to dry distillation. The final product is

(1) ethanal

(2) propanal

(3) propanone

(4) ethanol

55. The correct statement is

- BF₃ is the strongest Lewis acid among the other boron halides.
- (2) B1₃ is the weakest Lewis acid among the boron halides.
- (3) There is maximum pπ pπ back bonding in BF₃.
- (4) There is minimum pπ pπ back bonding in BF₃.

56. Which of the following compound possesses the "C - H" bond with the lowest bond dissociation energy?

(1) Toluene

(2) Benzene

(3) n-pentane

(4) 2, 2-dimethyl propane

- In presence of HCl, H₂S results the precipitation of Group-2 elements but not Gp-4 elements during qualitative analysis. It is due to
 - (1) higher concentration of S2-
- (2) higher concentration of H+
- (3) lower concentration of S²-
- (4) lower concentration of H+
- 58. One of the following conversion results in the change of hybridization and geometry:
 - (1) CH₄ to C₂H₆

(2) NH₃ to NH₄

(3) BF₃ to BF₄

- (4) H₂O to H₃O
- 59. Water softening by Clark's process uses
 - (1) CaHCO₁

(2) NaHCO₃

(3) Na₂CO₃

- (4) Ca(OH),
- 60. An alkali metal hydride (NaH) reacts with diborane in 'A' to give a tetrahedral compound 'B' which is extensively used as reducing agent in organic synthesis. The compounds 'A' and 'B' respectively are
 - (1) C₂H₆ and C₂H₅Na

(2) CH3COCH3 and B3N3H6

(3) CoHo and NaBH4

(4) (C2H5)2O and NaBH4





A-1

16

r